

# MARE'S MILK PLUST

## FOR NURSING, ORPHANED OR REJECTED FOALS

#### **GUARANTEED ANALYSIS**

Crude Protein, Min.	21.00%
Lysine, Min.	1.80%
Crude Fat, Min.	14.00%
Crude Fiber, Max.	0.15%
Calcium, Min.	0.70%
Calcium, Max.	1.20%
Phosphorus, Min.	0.65%
Copper, Min.	35 ppm
Zinc, Min.	110 ppm
Selenium, Min.	0.30 ppm
Vitamin A, Min.	12,000 IU/lb
Vitamin D3, Min.	1,200 IU/lb
Vitamin E, Min.	100 IU/lb
Ascorbic Acid, Min.	50 mg/lb

<sup>\*</sup>Not recognized by AAFCO as an essential nutrient.

## RECOMMENDED FEEDING PROGRAM FOR ORPHANED OR REJECTED FOALS

Age of Foal	Recommended Feed	
1-60 Days	Mare's Milk Plus™	
60-120 Days	Foal Starter™	
120+ Days	GRO 'N WIN™ or GROWTH	

For mares 1,000-1,500 pounds – one bag of Mare's Milk Plus™ will last 7-10 days.

### FEEDING DIRECTIONS

Colostrum, or first milk Contains high levels of antibodies and deeds to be consumed within the firs 12-24 hours. After colostrum has be ingested, it is safe to introduce the foal to Mare's Milk  $Plus^{M}$ .

FEEDING RATE			
Weight of Mare or Expected Mature Weight of Foal (LB)	Water (QT)	Mare's Milk Plus™ (Cups) Use enclosed measuring cup	
250	4	2	
500	8	4	
1,000	16	8	
1,500	24	12	
2,000	32	16	

Enclosed cup measures 272 grams of Mare's Milk Plus™

- ▶ Mare's Milk Plus™ should be mixed with cool tap water and stirred with a wire whisk.
- Mare's Milk Plus<sup>™</sup> should be fed at room temperature and does not require warming prior to ingestion.
- ▶ Start by feeding your foal half of the recommended volume from the table and gradually increase the amount over the next 10 days until the recommended amount is reached.

  Always discard any milk not consumed prior to adding fresh.
- Mare's Milk Plus<sup>™</sup> should be mixed every 12 hours; however, a 24-hour supply can be mixed if necessary. One half can be fed and the other half covered and stored in a cool place until the next feeding (does not require refrigeration).
- ▶ Always keep equipment clean to avoid the risk of infection.
- Provide clean, fresh water at all times.
- Doccasional diarrhea may occur in foals fed milk replacer. When this happens, reduce the amount of Mare's Milk Plus™ by 25% and then gradually increase until the foal is consuming the recommended amount. If diarrhea persists, contact your veterinarian.

#### **INGREDIENTS**

Dried Whey Protein Concentrate, Dried Whey, Dried Whey Product, Animal and Vegetable Fat (Preserved with BHA and BHT), Citric Acid, Potassium Sorbate (Preservative), L-Lysine Monohydrochloride, Dicalcium Phosphate, Calcium Carbonate, Lecithin, Vitamin A Acetate, Vitamin D3 Supplement, Vitamin E Supplement, Ascorbyl Phosphate, Choline Chloride, Folic Acid, Manganese Sulfate, Ferrous Sulfate, Copper Sulfate, Zinc Sulfate, Complex, Iron Polysaccharide Complex, Copper Polysaccharide Complex, Manganese Polysaccharide Complex, Cobalt Sulfate, Calcium Iodate, Sodium Selenite, Niacinamide, Calcium Pantothenate, Riboflavin Supplement, Vitamin B12 Supplement, Thiamine Mononitrate, Sodium Silico Aluminate, Mono and Diglycerides of Edible Fats or Oils, Artificial Flavor.

WARNING: CONTAINS ADDED COPPER. DO NOT FEED TO SHEEP

YOUR **PASSION.** OUR **PURPOSE.**<sup>TM</sup>