


**KEEP THIS CLIENT INFORMATION SHEET IN THE CARTON WITH THE SILEO DOSING SYRINGE
REVIEW PRIOR TO GIVING EACH DOSE TO YOUR DOG**

SILEO (dexmedetomidine oromucosal gel)

CLIENT INFORMATION SHEET FOR OWNER/HANDLER USE AND SAFETY

This sheet summarizes the basic information about SILEO and does not replace the instructions from your veterinarian. Talk to your veterinarian if you have questions regarding any part of this information or if you want to know more about SILEO. The syringe must be used in order to correctly dose the drug to your dog. You may use either a full, unused syringe or a partially used syringe that contains enough gel for the desired dose.



INSTRUCTIONS FOR USE

A. NEW SYRINGE SET UP BEFORE FIRST DOSING:

<p>WEAR GLOVES</p>	<p>1. Wear impermeable disposable gloves when handling SILEO and handling the oral syringe.</p>
<p>HOLD PLUNGER</p>	<p>2. Hold the oral syringe so that you can see the dot markings on the oral syringe plunger.</p>
<p>UNLOCK</p>	<p>3. Hold the plunger with your left hand and unlock the green ring-stop by turning it towards you until it is able to slide freely.</p>
<p>MOVE RING</p>	<p>4. Move the ring-stop to the opposite end of the plunger.</p>
<p>LOCK</p>	<p>5. Hold the plunger with your right hand and lock the ring-stop by turning it away from you. Confirm that the ring-stop is in the locked position on the dosing syringe by making sure it is not freely moveable.</p>

B. DOSE SELECTION AND DOSING:

<p>UNLOCK</p>	<p>6. Hold the plunger with your right hand and unlock the ring-stop by turning it towards you. Do not pull the plunger!</p>
<p>MOVE RING</p>	<p>7. Move the ring-stop towards the other end of the plunger for choosing the correct dose based on your veterinarian's prescription.</p>
<p>SET DOSE AND LOCK</p>	<p>8. Position the ring-stop so that the side nearest the barrel is in line with the graduation mark (black line), and the required number of dots shows between the ring-stop and the barrel. Lock the ring-stop by turning it away from you. Make sure that the ring-stop is in the locked position on the dosing syringe prior to administration in order to prevent accidental overdose.</p>
<p>PULL CAP (TIGHT)</p>	<p>9. Pull the cap strongly while holding the barrel. Note the cap is very tight (pull, do not twist). Save the cap for replacement. Confirm the ring-stop is still in the locked position on the dosing syringe by making sure that it is not freely moveable.</p>
<p>DOSE INTO CHEEK</p>	<p>10. Place the oral syringe tip between your dog's cheek and gum and press the plunger until the ring-stop causes the plunger to stop.</p>
<p>NOT SWALLOWED</p>	<p>IMPORTANT: The gel should not be swallowed. If the gel is swallowed, it may not be effective. Feeding and giving treats within 15 minutes after administration should be avoided.</p>
<p>BACK TO PACKAGE</p>	<p>11. Recap the oral syringe and return it to the outer package as the product is sensitive to light. Make sure that the carton is closed properly. Keep the package out of sight and reach of children at all times. Remove and discard gloves. Use or discard syringe contents within 4 weeks after opening the syringe.</p>