



Dog Owner Information about Carprofen Flavored Tablets for Osteoarthritis and Post-Surgical Pain

Generic name: carprofen ("car-prō-fen")

This summary contains important information about carprofen tablets. You should read this information before you start giving your dog carprofen tablets and review it each time the prescription is refilled. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or if you want to know more about carprofen tablets.

What are carprofen tablets?
Carprofen tablets are a nonsteroidal anti-inflammatory drug (NSAID) used to reduce pain and inflammation (swelling) due to osteoarthritis and pain following surgery in dogs. Carprofen tablets are a prescription drug for dogs. They are available as flavored tablets and are given to dogs by mouth.

Osteoarthritis (OA) is a painful condition caused by "wear and tear" of cartilage and other parts of the joints that may result in the following changes or signs in your dog:

- Limping or lameness
- Decreased activity or exercise (reluctance to stand, climb stairs, jump or run, or difficulty in performing these activities)
- Stiffness or decreased movement of joints

To control surgical pain (e.g. for surgeries such as spays, ear procedures or orthopedic repairs) your veterinarian may administer carprofen tablets before the procedure and recommend that your dog be treated for several days after going home.

What kind of results can I expect when my dog is on carprofen tablets?
While carprofen tablets are not a cure for osteoarthritis, they can relieve the pain and inflammation of OA and improve your dog's mobility.

- Response varies from dog to dog but can be quite dramatic.
- In most dogs, improvement can be seen in a matter of days.
- If carprofen tablets are discontinued or not given as directed, your dog's pain and inflammation may come back.

Who should not take carprofen tablets?
Your dog should not be given carprofen tablets if he/she:

- Has had an allergic reaction to carprofen, the active ingredient of carprofen tablets.
- Has had an allergic reaction to aspirin or other NSAIDs (for example deroxycob, etodolac, firocoxib, meloxicam, phenylbutazone or toposinai) such as hives, facial swelling, or red or itchy skin.

Carprofen tablets should be given to dogs only.
Cats should not be given carprofen tablets. Call your veterinarian immediately if your cat receives carprofen tablets. People should not take carprofen tablets. Keep carprofen tablets and all medicines out of reach of children. Call your physician immediately if you accidentally take carprofen tablets.

How to give carprofen tablets to your dog.
Carprofen tablets should be given according to your veterinarian's instructions. Your veterinarian will tell you what amount of carprofen tablets is right for your dog and for how long it should be given. Carprofen tablets should be given by mouth and may be given with or without food.

What to tell/ask your veterinarian before giving carprofen tablets.
Tell your veterinarian about:

- The signs of OA you have observed (for example limping, stiffness).
- The importance of weight control and exercise in the management of OA.
- What tests might be done before carprofen tablets are prescribed.
- How often your dog may need to be examined by your veterinarian.
- The risks and benefits of using carprofen tablets.

Tell your veterinarian if your dog has ever had the following medical problems:

- Experienced side effects from carprofen tablets or other NSAIDs, such as aspirin
- Digestive upset (vomiting and/or diarrhea)
- Liver disease
- Kidney disease
- A bleeding disorder (for example, Von Willebrand's disease)

Tell your veterinarian about:

- Any other medical problems or allergies that your dog has now or has had.
- All medicines that you are giving your dog or plan to give your dog, including those you can get without a prescription.

Tell your veterinarian if your dog is:

- Pregnant, nursing or if you plan to breed your dog.

What are the possible side effects that may occur in my dog during carprofen tablets therapy?

Carprofen tablets, like other drugs, may cause some side effects. Serious but rare side effects have been reported in dogs taking NSAIDs, including carprofen tablets. Serious side effects can occur with or without warning and in rare situations result in death. The most common NSAID-related side effects generally involve the stomach (such as bleeding ulcers), and liver or kidney problems. Look for the following side effects that can indicate your dog may be having a problem with carprofen tablets or may have another medical problem:

- Decreased or no increase in appetite
- Vomiting
- Change in bowel movements (such as diarrhea, or black, tarry or bloody stools)
- Change in behavior (such as decreased or increased activity level, incoordination, seizure or aggression)
- Yellowing of gums, skin, or whites of the eyes (jaundice)
- Change in drinking habits (frequency, amount consumed)
- Change in urination habits (frequency, color, or smell)
- Change in skin (redness, scabs, or scratching)

It is important to stop therapy and contact your veterinarian immediately if you think your dog has a medical problem or side effect from carprofen tablets therapy. If you have additional questions about possible side effects, talk to your veterinarian.

Can carprofen tablets be given with other medicines?
Carprofen tablets should not be given with other NSAIDs (for example, aspirin, deroxocob, etodolac, firocoxib, meloxicam, toposinai) or steroids (for example, cortisone, dexamethasone, prednisone, triamcinolone).

Tell your veterinarian about all medicines you are giving your dog in the past, and any medicines that you are planning to give with carprofen tablets. This should include other medicines that you can get without a prescription. Your veterinarian may want to check that all of your dog's medicines can be given together.

What do I do in case my dog eats more than the prescribed amount of carprofen tablets?
Contact your veterinarian immediately if your dog eats more than the prescribed amount of carprofen tablets.

What else should I know about carprofen tablets?
This sheet provides a summary of information about carprofen tablets. If you have any questions or concerns about carprofen tablets, or osteoarthritis, or postoperative pain, talk to your veterinarian.

As with all prescribed medicines, carprofen tablets should only be given to the dog for which it was prescribed. It should be given to your dog only for the condition for which it was prescribed. It is important to periodically discuss your dog's response to carprofen tablets at regular check ups. Your veterinarian will best determine if your dog is responding as expected and if your dog should continue receiving carprofen tablets.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet, contact Dechra at 1-866-933-2472. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-1088 or http://www.fda.gov/reportanimalae


Approved by FDA under ANADA # 200-681
Manufactured for:
Dechra Veterinary Products
7015 College Boulevard, Suite 525
Overland Park, KS 66211 USA
Rev. February 2020



Carprofen Tablets Flavored Tablets

Non-steroidal anti-inflammatory drug
For oral use in dogs only

CATION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
DESCRIPTION: Carprofen tablets (carprofen) are a non-steroidal anti-inflammatory drug (NSAID) of the propionic acid class that includes naproxen, ibuprofen, and ketorolac. Carprofen is the nonproprietary designation for a substituted carbocyclic 4-hydroxy-2-methyl-5H-cyclohex-2-ene-1-carboxylic acid. The empirical formula is C₁₇H₂₃NO₂ and the molecular weight 273.72. The chemical structure of carprofen is:



Carprofen is a white, crystalline compound. It is freely soluble in ethanol, but practically insoluble in water at 25°C.
CLINICAL PHARMACOLOGY: Carprofen is a non-steroidal anti-inflammatory agent with characteristic analgesic and anti-inflammatory activity approximately equivalent to indomethacin in animal models.¹ The mechanism of action of carprofen, like that of other NSAIDs, is believed to be associated with the inhibition of cyclooxygenase activity. Two unique cyclooxygenases have been described in mammals: the constitutive cyclooxygenase, COX-1, which is essential for normal gastrointestinal and renal function; the inducible cyclooxygenase, COX-2, generates prostaglandins necessary for normal gastrointestinal and renal function. The selective cyclooxygenase, COX-2, generates prostaglandins involved in inflammation. Inhibition of COX-1 is thought to be associated with gastrointestinal and renal toxicity while inhibition of COX-2 provides anti-inflammatory activity. The specificity of a particular NSAID for COX-2 versus COX-1 may vary from species to species.¹ In an *in vitro* study using canine cell cultures, carprofen demonstrated selective inhibition of COX-2 versus COX-1. Clinical relevance of these data has not been shown. Carprofen has also been shown to inhibit the release of several prostanoids in the inflammatory cell systems of polymorphonuclear leukocytes (PMN) and human mononuclear synovial cells, including inhibition of acute (PMN system) and chronic (synovial cell system) inflammatory mediators.²

Several studies have demonstrated that carprofen has modulatory effects on both humoral and cellular immune responses.^{3,4} Data also indicate that carprofen inhibits the production of osteoclast-activating factor (OAF), PGE₂, and TNF- α by its inhibitory effects on prostaglandin biosynthesis.⁵
Based upon comparison with data obtained from intravenous administration, carprofen is rapidly and nearly completely absorbed (over 90% bioavailable) when administered orally.⁶ Peak blood plasma concentrations are achieved in 1-3 hours after oral administration of 1, 5, and 25 mg/kg in dogs. The mean terminal half-life of carprofen is approximately 8 hours (range 4.5-8.8 hours) after single oral doses varying from 1-35 mg/kg of body weight. After a 100 mg single intravenous bolus dose, the mean elimination half-life was approximately 11.7 hours in the dog. Carprofen is more than 99% bound to plasma protein and exhibits a very small volume of distribution.

Carprofen is eliminated in the dog primarily by biotransformation in the liver followed by renal excretion of the resulting metabolites: the ester glucuronide of carprofen and the glucuronide of phenolic metabolites.⁷ Neither carprofen and its hydrolytic carprofen in the feces (70-80%) and urine (10-20%). Some enterogastric circulation of the drug is observed.

INDICATIONS: Carprofen tablets are indicated for the relief of pain and inflammation associated with osteoarthritis for the control of postoperative pain associated with soft tissue and orthopedic surgeries in dogs.
CONTRAINDICATIONS: Carprofen tablets should not be used in dogs exhibiting preexisting hypersensitivity to carprofen.

WARNINGS: Keep out of reach of children. Not for human use. Consult a physician in cases of accidental ingestion by humans. **For use in dogs only.** Do not use in cats.
All dogs should undergo a thorough history and physical examination before initiation of NSAID therapy. Appropriate laboratory tests to establish hematological and serum biochemical baseline data prior to, and periodically during, administration of any NSAID should be considered. **Owners should be advised to observe for signs of potential drug toxicity (see Information for Dog Owners, Adverse Reactions, Animal Safety and Post-Approval Experience).**

PRECAUTIONS: As a class, cyclooxygenase inhibitors (NSAIDs) may be associated with gastrointestinal, renal, and hepatic toxicity. Effects may result from decreased prostaglandin production and inhibition of the enzyme cyclooxygenase which is responsible for the formation of prostaglandins from arachidonic acid.⁸ While NSAIDs inhibit prostaglandins that cause inflammation, they may also inhibit those prostaglandins which maintain normal homeostatic functions. These anti-inflammatory effects may result in clinically significant disease in patients with underlying or pre-existing disease more often than in healthy patients.⁹ NSAID therapy could mask or obscure disease which has previously been undiagnosed due to the absence of apparent clinical signs. Patients with underlying renal disease, for example, may experience exacerbation or decompensation of their renal disease when NSAID therapy.¹⁰ The use of parenteral fluids during surgery should be considered to reduce the potential risk of renal complications when NSAIDs are administered.

Carprofen is an NSAID, and as with others of this class, adverse reactions may occur with its use. The most frequently reported effects have been gastrointestinal signs. Events involving suspected renal, hematologic, neurologic, dermatologic, and hepatic effects have also been reported. Patients at greatest risk for renal toxicity are those that are dehydrated, on concurrent diuretic therapy, or those with renal, cardiovascular, and/or hepatic dysfunction. Concurrent administration of potentially nephrotoxic drugs should be approached cautiously, with appropriate monitoring. Concurrent use of carprofen tablets with other anti-inflammatory drugs, such as other NSAIDs or corticosteroids, should be avoided because of the potential increase in adverse reactions, including gastrointestinal ulceration and perforation. Similarly, the drug-associated adverse reactions with the individual agent. Dogs that have experienced adverse reactions from one NSAID may experience adverse reactions from another NSAID. Carprofen tablets treatment was not associated with renal toxicity or gastrointestinal ulceration or renal colic safety studies of up to 10 times the dose of healthy dogs.

Carprofen tablets are not recommended for use in dogs with bleeding diatheses (e.g., von Willebrand's disease), as safety has not been established in dogs with these disorders. Safety can be safely use of carprofen tablets in animals with a variety of acute, pre-pregnant, dogs, dogs used for breeding purposes, or in lactating bitches has not been established. Studies to determine the safety of carprofen tablets when administered concomitantly with other pain-relief or analgesic medications have not been conducted. Drugs concomitantly should be monitored closely. In patients receiving additional therapy, such drugs commonly used include corticoids, anticonvulsant and behavioral medications. It has been suggested that treatment with carprofen may reduce the level of analgesic analgesics needed.¹¹

If additional pain medication is warranted after administration of the total daily dose of carprofen tablets, appropriate analgesia should be considered. The use of another NSAID is not recommended. Carprofen tablets approximate times when switching from one NSAID to another or when switching from conventional to NSAID use.

INFORMATION FOR DOG OWNERS: Carprofen tablets, like other drugs of this class, are not for human use. Owners should be advised of the potential for adverse reactions and of influence of the clinical signs associated with drug resistance. Adverse reactions may include decreased appetite, vomiting, diarrhea, dark or tarry stools, increased water consumption, increased urination, pain, gums due to anemia, yellowing of gums, skin, or whites of the eye due to jaundice. Intercourse, reproduction, sexual, or maternal care should be avoided during the treatment period. **Adverse reactions associated with this drug class can occur without warning and in rare situations result in death (see Adverse Reactions). Owners should be advised to discontinue carprofen tablets therapy and contact their veterinarian immediately if signs of intolerance are observed.** The great majority of patients with drug-related adverse reactions have recovered when the signs are recognized, the drug is withdrawn, and veterinary care, if appropriate, is initiated. Owners should be advised of the importance of periodic follow-up for all dogs during administration of any NSAID.

ADVERSE REACTIONS: During investigational studies of osteoarthritis with twice daily administration of 1 mg/kg, no clinically significant adverse reactions were reported. Some clinical signs were observed during field studies (n=397) which were similar for carprofen and placebo-treated dogs. Incidences of the following were observed in both groups: vomiting (1%), diarrhea (1%), changes in appetite (0%), lethargy (1.4%), behavioral changes (1%), and constipation (0.3%). The product vehicle served as control.

There were no serious adverse events reported during clinical field studies of osteoarthritis with once daily administration of 1 mg/kg. The following categories of abnormal health observations were reported. The product vehicle served as control.

Observation	Percentage of Dogs with Abnormal Health Observations Reported in Osteoarthritis Field Study (2 mg/kg once daily)	
	Carprofen Tablets (n=126)	Placebo (n=132)
Incidence	1.8	1.5
Vomiting	3.1	3.6
Diarrhea stool	3.2	4.5
Behavior change	0.8	0.8
Depression	0.8	0.5
PUPD	0.8	0.3
SAP increase	7.8	8.9
ALT increase	5.4	4.5
ACT increase	2.3	0.8
BUN increase	3.1	1.5
Bilirubin	0.2	0.1
Ketotaurin	14.7	9.1

Clinical pathology parameters listed represent reports of increases from pre-treatment values, medical judgment is necessary to determine clinical relevance.
During investigational studies of surgical pain for the tablet formulation, no clinically significant adverse reactions were reported. The product vehicle served as control.

Observation*	Percentage of Dogs with Abnormal Health Observations Reported in Surgical Pain Field Studies with Tablets (2 mg/kg once daily)	
	Carprofen Tablets (n=148)	Placebo (n=148)
Vomiting	10.1	13.1
Diarrhea stool	6.1	6.0
Incidence	2.3	1.0
Depression	1.4	0
Dysphytosis	0.7	0
ALT increase	1.4	0
BUN increase	0.7	0
Urea nitrogen	1.4	1.3
Wound drainage	1.4	0

* A single dog may have experienced more than one occurrence of an event.

Post-Approval Experience:
Although not all adverse reactions are reported, the following adverse reactions are based on voluntary post-approval adverse drug experience reporting. The categories of adverse reactions are listed in decreasing order of frequency by body system.

Systemic: Anemia, diarrhea, constipation, hypoproteinemia, melasma, hematemesis, gastrointestinal ulceration, gastrointestinal bleeding, anorexia.

Neurologic: Ataxia, paresis, paralysis, ataxia, vestibular signs, disorientation.

Uterine: Hemorrhage, pyometra, pyometra, urinary incontinence, urinary tract infection, azotemia, acute renal failure, leukocytosis, leukopenia, leukopenia, hypocalcemia, hypoproteinemia. Approximately one fourth of hepatic signs were in a Labrador Retriever.

Behavioral: Sedation, lethargy, hypersensitivity, restlessness, aggressiveness.

Hematologic: Immune-mediated hemolytic anemia, immune-mediated thrombocytopenia, blood loss anemia, epistaxis.

Cardiologic: Bradycardia, decreased atrioventricular conduction, tachycardia; most dramatic (flat axis), nonrecording, paroxysmal tachycardia, ventricular tachycardia.

Immunologic or hypersensitivity: Facial swelling, hives, erythema.

In rare situations, death has been associated with some of the adverse reactions listed above. To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet, contact Dechra at 1-866-933-2472. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-1088 or http://www.fda.gov/reportanimalae

DOSEAGE AND ADMINISTRATION: Always provide Client Information Sheet with prescription. Carefully consider the potential benefits and risk of carprofen tablets and other treatment options before deciding to use carprofen tablets. Use the lowest effective dose for the shortest duration consistent with individual response. The recommended dosage for oral administration to dogs is 2 mg/kg (4 mg/kg of body weight) daily, may be administered as 2 mg/kg of body weight once daily divided and administered as 1 mg/kg (2 mg/kg twice daily) for the control of postoperative pain, administered approximately 2 hours before the procedure. Tablets may score and dosage should be calculated to half-tablet increments.

EFFECTIVENESS: Continuation of the effectiveness of carprofen tablets for the relief of pain and inflammation associated with osteoarthritis, and for the control of postoperative pain associated with soft tissue and orthopedic surgeries was demonstrated in a placebo-controlled, randomised studies examining the anti-inflammatory and analgesic effectiveness of carprofen tablets in various breeds of dogs.

Separate placebo-controlled, masked, multicenter field studies confirmed the anti-inflammatory and analgesic effectiveness of carprofen tablets when dosed at 2 mg/kg once daily or when divided and administered at 1 mg/kg twice daily. In these low field studies, dogs dosed with carprofen tablets showed statistically significant weight improvement based on farmowner evaluations by the veterinarian and owner observations when administered carprofen tablets compared to placebo.

Separate double-blind, randomized, multicenter field studies confirmed the effectiveness of carprofen tablets for the control of postoperative pain when dosed at 2 mg/kg once daily in various breeds of dogs. In these studies, dogs presented for orthopedic/neurologic repair and/or surgery were administered carprofen tablets preoperatively and for a maximum of 7 days (with 6 days of drug) postoperatively. In general, dogs administered carprofen tablets showed statistically significant improvement in pain scores compared to controls.

ADVERSE EFFECTS: Laboratory studies in various breeds of dogs and clinical field studies have demonstrated that carprofen tablets are well tolerated in dogs after oral administration.

In a large animal safety study, carprofen tablets were administered orally to healthy Beagle dogs at 1, 5, and 10 mg/kg twice daily for 7, 14, and 28 days. The recommended total daily dose is 42 mg/kg once daily. In these low field studies, dogs dosed with carprofen tablets showed statistically significant weight improvement based on farmowner evaluations by the veterinarian and owner observations when administered carprofen tablets compared to placebo.

In a 28-week study, minor dermatologic changes occurred in dogs in each of the treatment groups but not in the control group. The changes were described as slight redness or dry skin and were diagnosed as non-specific dermatitis. The probability exists that these mild lesions were treatment related, but no dose relationship was observed.

Clinical field studies were conducted with dogs of different breeds at the recommended dose and for 14 days (597 dogs were included in a study evaluating 1 mg/kg twice daily and 250 dogs were included in separate study evaluating 2 mg/kg once daily). In both studies the dogs were clinically well tolerated and the incidence of clinical adverse reactions to carprofen tablets treated animals was higher than placebo-treated animals. Observed adverse reactions included hives in carprofen tablets. For animals receiving 1 mg/kg twice daily, the mean post-treatment serum alanine aminotransferase (ALT) was 9.1 IU/l (range 0.1-24.4 IU/l) after treatment with 1 mg/kg twice daily carprofen tablets and placebo, respectively. Differences were not statistically significant. For animals receiving 2 mg/kg once daily, the mean post-treatment serum ALT was 4.3 IU/l (range 0.1-16.0 IU/l) after treatment with 2 mg/kg once daily carprofen tablets and placebo, respectively. For animals receiving 1 mg/kg twice daily, the mean post-treatment serum ALT was 6.4 IU/l (range 0.1-24.4 IU/l) after treatment with 1 mg/kg twice daily carprofen tablets and placebo, respectively. Changes in the clinical laboratory values (hematology and clinical chemistry) indices of hematopoietic, renal, hepatic, and coagulation function were not clinically significant. The mean post-treatment ALT values were 7.1 IU/l and 2.5 IU/l for the pre-treatment values for dogs receiving carprofen tablets and placebo, respectively. The mean post-treatment ALT values were 3.1 IU/l for the pre-treatment values for dogs receiving placebo.

STORAGE: Store at 20°C to 25°C (68°F to 77°F).

HOW SUPPLIED: Carprofen flavored tablets are scored, and contain 25 mg, 75 mg, and 100 mg of carprofen per tablet. Each tablet size is packaged in bottles containing 10, 30, and 100 tablets.

Carprofen flavored tablets 25 mg, 100 tablets NDC 17033-422-30
Carprofen flavored tablets 75 mg, 60 tablets NDC 17033-422-60
Carprofen flavored tablets 100 mg, 180 tablets NDC 17033-422-180
Carprofen flavored tablets 25 mg, 30 tablets NDC 17033-422-30
Carprofen flavored tablets 75 mg, 60 tablets NDC 17033-422-60
Carprofen flavored tablets 100 mg, 180 tablets NDC 17033-422-180

REFERENCES:
1. Smith, J., et al. Anti-inflammatory and Anti-Rheumatic Drugs, Vol. 4. New and Anticancer Drugs, 949-951, ed. CRC Press, Boca Raton, pp. 33-47, 1986.

2. von, J., Buring, H.E. Mechanism of action of anti-inflammatory drugs. *Scand J Rheumatism* 22:102, pp. 9-21, 1993.

3. Grossman, C.J., Williams, J., Lucas, P.S., et al. Inhibition of constitutive and inducible cyclooxygenase activity in human platelets and mononuclear cells by NSAIDs and COX-2 inhibitors. *Inflammation Research* 44:253-257, 1995.

4. Nishida, H., Uemura, M., Saitoh, S.P. Evaluation of selective inhibition of cyclooxygenase-1 and 2 by carprofen and other nonsteroidal anti-inflammatory drugs. *Am J Vet Res* 58:11, pp. 1441-1446, November 1998.

5. Croppers, J., et al. Endogenous prostaglandin E₂ enhances polynuclear immunoglobulin production by orally inhibiting T suppressor cell activity. *Cell Immunol* 79:41, 1982.

7. Schenker, P.P., et al. Effect of prostaglandin synthetase inhibition on the immune response. *Immunopharmacology* 3:205, 1981.

8. Long, M., et al. Modulation of the development of cell mediated immunity: possible roles of the products of cyclooxygenase and lipooxygenase pathways of arachidonic acid metabolism. *Int J Immunopharmacology* 16: 182, 1993.

9. Wet, S.C. Immunoregulatory activity of cultured-induced suppressor macrophages. *Cell Immunol* 72:14, 1982.

10. Scovetti, A., et al. Biopharmaceutical evaluation of carprofen following single intravenous, oral, and rectal doses in dogs. *Biopharm Drug Dispos* 11(2):185-194, 1990.

11. New AM. Toxicology of nonsteroidal anti-inflammatory drugs. *Veterinary Clinics of North America, Small Animal Practice* 20, March 1990.

12. New AM. Pathogenesis and pathophysiology of ischemic injury in cases of acute renal failure. *Compend for Cont Ed* 18:1, January 1994.

13. Bettle DM. Prostaglandins: Physiology and clinical implications. *Compend for Cont Ed* 11, November 1984.

14. Saitoh H. Nonsteroidal anti-inflammatory drugs, prostaglandins, and the kidney. *JAMA* 183:8, May 1980.

15. Ke, C.H., Lange, O.H., Mandagaj, R.E., et al. Effects of subtherapeutic and carprofen on the minimal alveolar concentration of halothane in dogs. *AAEP* 49:1702-1703, 2000.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet, contact Dechra at 1-866-933-2472. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-1088 or http://www.fda.gov/reportanimalae

Approved by FDA under NDACA # 200-681
Manufactured for:
Dechra Veterinary Products
7015 College Boulevard, Suite 525
Overland Park, KS 66211 USA
Rev. February 2020



Product:.....Carprofen Tablets Flavored - US - Leaflet	Proof:.....	Date:.....	Proof:.....	Date:.....
Dimensions:.....7.75" x 18"	3.1 (L)	12-02-2020		
Primary brand name font size:.....21pt - (Dog owner 24pt)	3.2 (L)	13-02-2020		
Established name.....17.85pt - (Dog owner 20.4pt)	3.3 (L)	18-02-2020		
Primary brand description font size:.....12.6pt - (Dog owner 14.4pt)	3.4 (L)	30-03-2020		
Body text font size:.....6pt - (Dog owner 9pt)	3.5 (L)	30-03-2020		
Item code:.....Rev. February 2020	3.6 (L)	01-04-2020		
Pharmacode:.....N/A				

Pantone reference guide
Colours to be printed:
BLACK

Do not print
CUTTER GUIDE TEXT AREA
CREASE CUT MARKS
GLUE PANEL REGISTRATION MARKS

STYLE DEVIATIONS
Colors to be printed:
Header - ("car-pro-fen") has had to use the "World" version of Helvetica owing to the accent above the "v".

REGULATORY AUTHORITIES' REQUESTS

Dechra