

FRONTLINE® SHIELD FOR DOGS

For use **ONLY** on dogs 9 weeks or older and weighing 41-80 lbs.

ACTIVE INGREDIENTS

Fipronil	6.01%
Permethrin*	44.88%
Pyriproxyfen	1.8%
OTHER INGREDIENTS	47.31%
TOTAL	100.0%

*cis/trans ratio: max 55% (±) cis and min 45% (±) trans



KEEP OUT OF REACH OF CHILDREN WARNING

For additional Precautionary Statements, refer to carton back panel. For additional Directions for Use and First Aid information, refer to carton inside panel.

FRONTLINE® SHIELD FOR DOGS

Convenient, long-lasting monthly topical for treatment of fleas, ticks, and chewing lice, and repellency of mosquitoes and stable flies. Effectively controls existing infestations and prevents re-infestations of fleas, ticks and chewing lice.

Ticks:

- Starts killing ticks in 1 hour, starting 2 days after treatment
- Repels and kills ticks for an entire month, including Deer ticks, Brown Dog ticks, American Dog ticks, and Lone Star ticks
- Kills ticks which may transmit Lyme disease, Rocky Mountain spotted fever, Anaplasmosis, Babesiosis, canine Ehrlichiosis or cause tick paralysis

Fleas:

- Starts killing fleas in 5 minutes, starting 2 days after treatment
- Kills fleas, flea eggs and flea larvae for an entire month
- Prevents development of flea eggs and flea larvae, effectively breaking the flea life-cycle
- Kills fleas which may cause flea allergy dermatitis and that may transmit bartonellosis, tularemia and tapeworm

Mosquitoes and Flies:

- Repels mosquitoes that may transmit heartworm disease and West Nile virus
- Repels and reduces blood-feeding by stable flies and mosquitoes for a month

Lice and Mites:

- Kills chewing lice
- Aids in control of Sarcoptes mite infestations which may cause sarcoptic mange

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. **FOR EXTERNAL USE ON DOGS ONLY. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL AND DIRECTIONS BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.**

Apply **FRONTLINE® SHIELD FOR DOGS** as follows:



1 TO REMOVE

Applicator:
Prior to opening **separate one blister** by tearing along perforation.

2

TO OPEN, use scissors to cut along dotted line of single blister, then pull apart foil and remove the applicator. **OR** fold on line at corner, **tear and open at slit** and remove the applicator.



3 HOLD Applicator Upright

away from you and your dog's face, cut applicator tip with scissors to open. Do not get product in your dog's eyes or mouth.



4 APPLY FRONTLINE® SHIELD FOR DOGS

Part dog's hair until skin is visible. Place applicator tip directly against exposed skin. Apply the product evenly to three spots on the dog's back, starting between the shoulder blades and continuing on to the second and third spots, squeezing the applicator until empty. Ensure that **FRONTLINE® SHIELD FOR DOGS** is not applied superficially on dog's hair.

FREQUENCY OF APPLICATION

Apply **FRONTLINE® SHIELD FOR DOGS** monthly to start killing ticks in 1 hour and fleas in just 5 minutes beginning 2 days after treatment. **FRONTLINE® SHIELD FOR DOGS** controls fleas, flea eggs, and flea larvae, ticks, and chewing lice infestations. Ticks, stable flies and mosquitoes are repelled for one month by a single application of **FRONTLINE® SHIELD FOR DOGS**.

FRONTLINE® SHIELD FOR DOGS remains effective even after bathing, water immersion, or exposure to sunlight 24 hours after application. A monthly application of **FRONTLINE® SHIELD FOR DOGS** kills existing fleas and prevents new fleas from developing; kills all stages of ticks, chewing lice, and repels mosquitoes and stable flies.

RESTRICTIONS

Do not allow children to apply product. Do not allow adults, children, or other pets to touch treated area until dry.

Do not use on cats or rabbits; or any animals other than dogs. Do not allow your dog to ingest this product. Do not use on dogs under 9 weeks of age or weighing less than 41 lbs. Do not apply more than once a month, or use more than one applicator per dog, regardless of the dog's weight. Do not split applicators between dogs. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Overdosing your dog can result in serious illness.

Do not bathe your dog within the first 24 hours after the product has been applied. Remember, your dog needs to be dry at the time **FRONTLINE® SHIELD FOR DOGS** is applied and remain dry 24 hours after application. If your dog is exhibiting signs of and/or is being treated for skin problems, talk to your veterinarian before applying any topical flea and tick control product. For use **ONLY** on dogs 9 weeks or older and weighing 41-80 lbs. to kill fleas and ticks and prevent fleas from developing; to repel and reduce blood-feeding by ticks, mosquitoes and stable flies; to kill chewing lice and to aid in control of Sarcoptes mites. Apply the entire contents of the applicator at each treatment. It may be helpful to remove the collar before application, so it doesn't absorb the product.

PESTICIDE STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store unused product in original container only in a cool dry place, out of reach of children and animals.

PESTICIDE DISPOSAL and CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and handling of this material when such use and handling are contrary to the label instructions.

Questions or Comments?

Call: 1-888-637-4251

EPA Reg. No. 65331-9
Boehringer Ingelheim Animal Health USA Inc.
3239 Satellite Blvd., Duluth, GA 30096

FRONTLINE® is a registered trademark of the Boehringer Ingelheim Group.
©2020 Boehringer Ingelheim Animal Health USA Inc. All rights reserved.

